



SWATH®技术发表文章目录



主要内容

SWATH®技术发表文章目录

蛋白组学

► 一、方法学	3.
► 二、翻译后修饰蛋白组学	6.
► 三、疾病蛋白组学	8.
► 四、植物蛋白组学	14.
► 五、动物蛋白组学	15.
► 六、微生物蛋白组学(发酵)	16.
► 七、蛋白相互作用、食品组学、生命科学研究等蛋白组学应用	18.
代谢组学	21.
脂质组学	23.
药物分析	24.
毒物、污染物分析	25.

蛋白组学

一、方法学

1. Open Swath enables automated, targetd analysis of data-independent acquisition MS data. Nat Biotechnol 2014, DOI:10.1038/nbt.2841, 影响因子：41.667
2. a multicenter study benchmarks software tools for label-free proteome quantification. Nat Biotechnol 2016, doi:10.1038/nbt.3685, 影响因子：41.667
3. TRIC: an automated alignment strategy for reproducible protein quantification in targeted proteomics. Nat Med 2016, doi: 10.1038/nmeth.3954 , 影响因子：32.621
4. DIA-Umpire: comprehensive computational framework for data independent acquisition proteomics. Nat Med 2015, doi:10.1038/nmeth.3255, 影响因子：32.621
5. Specter: linear deconvolution for targeted analysis of data-independent acquisition mass spectrometry proteomics. Nature method 2018, doi. org/10.1038/nmeth.4643 , 影响因子：29.886
6. Building ProteomeTools based on a complete synthetic human proteome. Nature method 2017, doi:10.1038/nmeth.4153 , 影响因子：29.886.
7. Building high-quality assay libraries for targeted analysis of SWATH MS data. Nature protocol 2015, doi:10.1038/nprot.2015.015 , 影响因子：12.423
8. Multi-laboratory assessment of reproducibility, qualitative and quantitative performance of SWATH-mass spectrometry. Nat Commun 2017, doi: 10.1038/s41467-017-00249-5, 影响因子：12.124
9. Complex-centric proteome profiling by SEC-SWATH-MS. Mol Syst Biol

2019, DOI 10.15252/msb.20188438, 影响因子: 10.581

10. Expansion of the Ion Library for Mining SWATH-MS Data through Fractionation Proteomics. *Anal Chem* 2014, doi:10.1021/ac501828a, 影响因子: 6.32
11. A reference peptide database for proteome quantification based on experimental mass spectrum response curves. *Bioinformatics* 2018, DOI: 10.1093/bioinformatics/bty201, 影响因子: 5.481
12. Targeted data extraction of the MS/MS spectra generated by data independent acquisition:a new concept for consistent and accurate proteome analysis. *Mol Cell Proteomics* 2012, O111.016717, 影响因子: 5.232
13. MS1 Peptide Ion Intensity Chromatograms in MS2 (SWATH) Data Independent Acquisitions. Improving Post Acquisition Analysis of Proteomic Experiments. *Mol Cell Proteomics* 2015, O115.048181, 影响因子: 5.232
14. Online Peptide Fractionation Using a Multiphasic Microfluidic Liquid Chromatography Chip Improves Reproducibility and Detection Limits for Quantitation in Discovery and Targeted Proteomics. *Mol Cell Proteomics* 2015, doi: 10.1074/mcp.M114.046425, 影响因子: 5.232
15. SWATH mass spectrometry performance using extended peptide MS/MS assay libraries. *Mol Cell Proteomics* 2016, doi: 10.1074/mcp.M115.055558, 影响因子: 5.232
16. Conserved peptide fragmentation as a benchmarking tool for mass spectrometers and a discriminating feature for targeted proteomics. *Mol Cell Proteomics* 2014, doi: 10.1074/mcp.O113.036475, 影响因子: 5.232
17. Numerical compression schemes for proteomics mass spectrometry data. *Mol Cell Proteomics* 2014, doi: 10.1074/mcp.O114.037879, 影响因子: 5.232
18. mzDB: a file format using multiple indexing strategies for the efficient analysis of large LC-MS/MS and SWATH-MS datasets. *Mol Cell Proteomics*

2015, doi: 10.1074/mcp.O114.039115, 影响因子: 5.232

19. Automated Validation of Results and Removal of Fragment Ion Interferences in Targeted Analysis of Data Independent Acquisition MS using SWATH-Prophet. *Mol Cell Proteomics* 2015, doi: 10.1074/mcp.O114.044917, 影响因子: 5.232
20. MS1 Peptide Ion Intensity Chromatograms in MS2 (SWATH) Data Independent Acquisitions. Improving Post Acquisition Analysis of Proteomic Experiments. *Mol Cell Proteomics* 2015, doi: 10.1074/mcp.O115.048181, 影响因子: 5.232
21. Targeted Data Extraction of the MS/MS Spectra Generated by Data-independent Acquisition: A New Concept for Consistent and Accurate Proteome Analysis. *Mol Cell Proteomics* 2012, DOI: 10.1074/mcp.O111.016717, 影响因子: 5.232
22. Identification of a Set of Conserved Eukaryotic Internal Retention Time Standards for Data-Independent Acquisition Mass Spectrometry. *Mol Cell Proteomics* 2015, mcp.O114.042267, 影响因子: 5.232
23. Effective leveraging of targeted search spaces for improving peptide identification in MS/MS based proteomics. *J Proteome Res* 2015, doi:10.1021/acs.jproteome.5b00504, 影响因子: 3.95
24. Pseudo isobaric peptide termini labelling for relative proteome quantification by SWATH MS acquisition. *Analyst* 2016, DOI: 10.1039/c6an00388e, 影响因子: 3.864
25. SWATH enables precise label-free quantification on proteome-scale. *Proteomics* 2015, doi:10.1002/pmic.201400270, 影响因子: 3.532
26. Automated measurement of site-specific N-glycosylation occupancy with SWATH-MS. *Proteomics* 2015, doi:10.1002/pmic.201400465, 影响因子: 3.532
27. Minimal sample requirement for highly multiplexed protein quantifica-

- tion in cell lines and tissues by PCT-SWATH mass spectrometry. *Proteomics* 2015, doi:10.1002/pmic.201500161, 影响因子：3.532
- 28. Effect of Peptide Assay Library Size and Composition in Targeted Data Independent Acquisition-Mass Spectrometry Analyses. *Proteomics* 2016, doi: 10.1002/pmic.201600007, 影响因子：3.532
 - 29. Processing strategies and software solutions for data-independent acquisition in mass spectrometry. *Proteomics* 2015, DOI:10.1002/pmic.201400323, 影响因子：3.532
 - 30. A first dataset toward a standardized community-driven global mapping of the human immunopeptidome. *Data in Brief* 2016, doi:10.1016/j.dib.2016.02.016, 影响因子：0.341
 - 31. Methods for SWATH®: Data Independent Acquisition on TripleTOF Mass Spectrometers. *Methods Mol Biol* 2016, DOI 10.1007/978-1-4939-3524-6_16
 - 32. The beauty of being (label)-free: sample preparation methods for SWATH-MS and next-generation targeted proteomics. *F1000 Research* 2013, doi: 10.12688/f1000research.2-272
 - 33. Automated SWATH Data Analysis Using Targeted Extraction of Ion Chromatograms. *Methods Mol Biol* 2017, doi: 10.1007/978-1-4939-6747-6_20

二、翻译后修饰蛋白组学

- 1. Mass Spectral Enhanced Detection of Ubis Using SWATH Acquisition: MEDUSA-Simultaneous Quantification of SUMO and Ubiquitin-Derived Isopeptides. *J Am Chem Soc* 2014, DOI: 10.1007/s13361-014-0835-x, 影响因子：13.858
- 2. Similarities and Differences of Blood N-Glycoproteins in Five Solid Carcinomas at Localized Clinical Stage Analyzed by SWATH-MS. *Cell Reports* 2018,

doi:10.1016/j.celrep.2018.04.114, 影响因子: 8.282

3. SWATH® Analysis for Characterization and Quantification of Histone Post-translational Modifications. *Mol Cell Proteomics* 2015, O114.046102, 影响因子: 5.232
4. Sequential Window Acquisition of all Theoretical Mass Spectra (SWATH) Analysis for Characterization and Quantification of Histone Post-translational Modifications. *Mol Cell Proteomics* 2015, doi: 10.1074/mcp.O114.046102, 影响因子: 5.232
5. Phosphoprotein secretome of tumor cells as a source of candidates for breast cancer biomarkers in plasma. *Mol Cell Proteomics* 2014, doi: 10.1074/mcp.M113.035485, 影响因子: 5.232
6. Development of diagnostic fragment ion library for glycated peptides of Human serum albumin: Targeted quantification in prediabetic, diabetic and microalbuminuria plasma by PRM, SWATH and MSE. *Mol Cell Proteomics* 2015, doi: 10.1074/mcp.M115.050518, 影响因子: 5.232
7. Glycation inhibitors extend yeast chronological lifespan by reducing advanced glycation end products and by back regulation of proteins involved in mitochondrial respiration. *J Proteomics* 2017, doi:10.1016/j.jprot.2017.01.015, 影响因子: 3.722
8. Automated measurement of site-specific N-glycosylation occupancy with SWATH-MS. *Proteomics* 2015, doi:10.1002/pmic.201400465, 影响因子: 3.532
9. Quantitative measurements of N-linked glycoproteins in human plasma by SWATH-MS. *Proteomics* 2013, doi: 10.1002/pmic.201200417, 影响因子: 3.532
10. Label-free quantitative phosphoproteomics reveals regulation of vaso-dilator-stimulated phosphoprotein upon stathmin-1 silencing in a pair of isogenic colorectal cancer cell lines. *Proteomics* 2018, doi: 10.1002/

pmic.201700242, 影响因子: 3.532

11. Selective phosphorylation during early macrophage differentiation. Proteomics 2015, DOI: 10.1002/pmic.201400511, 影响因子: 3.532
12. Quantification of Lysine Acetylation and Succinylation Stoichiometry in Proteins Using Mass Spectrometric Data-Independent Acquisitions (SWATH). J Am Soc Mass Spect 2016, DOI:10. 1007/s13361-016-1476-z, 影响因子: 2.869

三、疾病蛋白组学

1. Inference and quantification of peptidoforms in large sample cohorts by SWATH-MS. Nat Biotechnol 2017, doi: 10.1038/nbt.3908, 影响因子: 41.667
2. Systems proteomics of liver mitochondria function. Science 2016, DOI: 10.1126/science.aad0189, 影响因子: 41.058
3. Rapid mass spectrometric conversion of tissue biopsy samples into permanent quantitative digital proteome maps. Nat Med 2015, doi:10.1038/nm.3807, 影响因子: 32.621
4. Absolute Proteome Composition and Dynamics during Dormancy and Resuscitation of Mycobacterium tuberculosis. Cell Host Microbe 2015, doi:10.1016/j.chom.2015.06.001, 影响因子: 14.946
5. Systematic proteome and proteostasis profiling in human Trisomy 21 fibroblast cells. Nat Commun 2017, DOI: 10.1038/s41467-017-01422-6, 影响因子: 12.124
6. Quantitative variability of 342 plasma proteins in a human twin population. Mol Syst Biol 2015, DOI: 10.15252/msb.20145728, 影响因子: 10.581
7. Similarities and Differences of Blood N-Glycoproteins in Five Solid Carcinomas at Localized Clinical Stage Analyzed by SWATH-MS. Cell Reports 2018,

- doi:10.1016/j.celrep.2018.04.114, 影响因子: 8.282
- 8. Impact of Alternative Splicing on the Human Proteome. *Cell Reports* 2017, doi:10.1016/j.celrep.2017.07.025, 影响因子: 8.282
 - 9. An open-source computational and data resource to analyze digital maps of immunopeptidomes. *eLife* 2015, doi: 10.7554/eLife.07661, 影响因子: 7.616
 - 10. Cell Autonomous and Noncell-Autonomous Role of NF- κ B p50 in Astrocyte-Mediated Fate Specification of Adult Neural Progenitor Cells. *glia* 2016, doi:10.1002/glia.23085, 影响因子: 5.846
 - 11. A Rapid Extraction Method for Glycogen from Formalin-fixed Liver. Carbohydrate Polymers 2015, doi:10.1016/j.carbpol.2014.11.005, 影响因子: 5.518
 - 12. A repository of assays to quantify 10,000 human proteins by SWATH-MS. *Scientific Data* 2014, DOI: 10.1038/sdata.2014.31, 影响因子: 5.305
 - 13. Data Descriptor: A tissue-based draft map of the murine MHC class I immunopeptidome. *Scientific Data* 2018, DOI: 10.1038/sdata.2018.157, 影响因子: 5.305
 - 14. Glycoproteomic Analysis of Prostate Cancer Tissues by SWATH Mass Spectrometry Discovers N-acylethanolamine Acid Amidase and Protein Tyrosine Kinase 7 as Signatures for Tumor Aggressiveness. *Mol Cell Proteomics* 2014, doi:10.1074/mcp.M114.038273, 影响因子: 5.232
 - 15. Glycoproteomic analysis of prostate cancer tissues by SWATH mass spectrometry. *Mol Cell Proteomics* 2014, doi: 10.1074/mcp.M114.038273, 影响因子: 5.232
 - 16. Phosphoprotein secretome of tumor cells as a source of candidates for breast cancer biomarkers in plasma. *Mol Cell Proteomics* 2014, doi: 10.1074/mcp.M113.035485, 影响因子: 5.232

17. Development of diagnostic fragment ion library for glycated peptides of Human serum albumin: Targeted quantification in prediabetic, diabetic and microalbuminuria plasma by PRM, SWATH and MSE. Mol Cell Proteomics 2015, doi: 10.1074/mcp.M115.050518, 影响因子: 5.232
18. In vivo and in vitro proteome analysis of HIV-1 infected, human CD4+ T cells. Mol Cell Proteomics 2017, M116.065235, 影响因子: 5.232
19. Label-free quantification of differentially expressed proteins in mouse liver cancer cells with high and low metastasis rates by a SWATH acquisition method. Sci Chin Chem 2014, DOI :10.1007/s11426-014-5093-z, 影响因子: 4.448
20. Proteomic profiling of HIV-infected T-cells by SWATH mass spectrometry. Virology 2018, doi:10.1016/j.virol.2018.01.025 , 影响因子: 4.368
21. Understanding the dimorphic lifestyles of human gastric pathogen Helicobacter pylori using the SWATH-based proteomics approach. Sci Rep 2016, DOI: 10.1038/srep26784, 影响因子: 4.122
22. Proteomics of thyroid tumours provides new insights into their molecular composition and changes associated with malignancy. Sci Rep 2016, DOI: 10.1038/srep23660, 影响因子: 4.122
23. SWATH-based proteomics identified carbonic anhydrase 2 as a potential diagnosis biomarker for nasopharyngeal carcinoma. Sci Rep 2017, DOI: 10.1038/srep41191, 影响因子: 4.122
24. Quantitative proteomics by SWATH-MS reveals sophisticated metabolic reprogramming in hepatocellular carcinoma tissues. Sci Rep 2017, DOI: 10.1038/srep45913, 影响因子: 4.122
25. Biomarker Discovery and Verification of Esophageal Squamous Cell Carcinoma Using Integration of SWATH/MRM. J Proteome Res 2015, doi: 10.1021/acs.jproteome.5b00438, 影响因子: 3.95
26. Osteogenic Commitment and Differentiation of Human Mesenchymal Stem

Cells by Low-Intensity pulsed Ultrasound Stimulation. *J Cell Physiol* 2017, doi:10.1002/jcp.26058, 影响因子：3.923

27. SWATH®- and iTRAQ-based quantitative proteomic analyses reveal an over-expression and biological relevance of CD109 in advanced NSCLC. *J Proteomics* 2014, doi:10.1016/j.jprot.2014.03.007, 影响因子：3.722
28. The secretome signature of malignant mesothelioma cell lines. *J Proteomics* 2016, doi:10.1016/j.jprot.2016.02.021, 影响因子：3.722
29. Multipronged quantitative proteomics reveals serum proteome alterations in breast cancer intrinsic subtypes. *J Proteomics* 2017, doi: 10.1016/j.jprot.2017.05.007, 影响因子：3.722
30. SWATH analysis of the synaptic proteome in Alzheimer's disease. *Neurochem Int* 2015, doi:10.1016/j.neuint.2015.04.004, 影响因子：3.603
31. Automated measurement of site-specific N-glycosylation occupancy with SWATH-MS. *Proteomics* 2015, doi: 10.1002/pmic.201400465, 影响因子：3.532
32. Quantitative measurements of N-linked glycoproteins in human plasma by SWATH-MS. *Proteomics* 2013, doi:10.1002/pmic.201200417, 影响因子：3.532
33. Label-free quantitative phosphoproteomics reveals regulation of vaso-dilator-stimulated phosphoprotein upon stathmin-1 silencing in a pair of isogenic colorectal cancer cell lines. *Proteomics* 2018, doi:10.1002/pmic.201700242, 影响因子：3.532
34. Two methods for proteomic analysis of formalin-fixed, paraffin embedded tissue result in differential protein identification, data quality, and cost. *Proteomics* 2015, doi:10.1002/pmic.201500147, 影响因子：3.532
35. Unravelling the proteome of degenerative human mitral Valves. *Proteomics* 2015, doi:10.1002/pmic.201500040, 影响因子：3.532

36. Selective phosphorylation during early macrophage differentiation. *Proteomics* 2015, DOI: 10.1002/pmic.201400511, 影响因子: 3.532
37. Identification of protein abundance changes in hepatocellular carcinoma tissues using PCT-SWATH. *Proteomics* 2018, doi:10.1002/prca.201700179, 影响因子: 3.532
38. Unveiling the Differences of Secretome of Human Bone Marrow Mesenchymal Stem Cells, Adipose Tissue-Derived Stem Cells, and Human Umbilical Cord Perivascular Cells: A Proteomic Analysis. *STEM CELLS DEVELOP* 2016, DOI: 10.1089/scd.2016.0048, 影响因子: 3.315
39. Applying SWATH Mass Spectrometry to Investigate Human Cervicovaginal Fluid During the Menstrual Cycle. *Biol Reprod* 2015, DOI: 10.1095/biolreprod.115.128231, 影响因子: 3.184
40. Quantitative analysis of hydroxyapatite-binding plasma proteins in genotyped individuals with late-stage age-related macular degeneration. *Exp Eye Res* 2017, doi:10.1016/j.exer.2018.03.023, 影响因子: 3.152
41. MVP immunohistochemistry is a useful adjunct in distinguishing leiomyosarcoma from leiomyoma and leiomyoma with bizarre nuclei. *Human Pathology* 2018, doi:10.1016/j.humpath.2017.12.020, 影响因子: 3.125
42. Using Data-Independent, High Resolution Mass Spectrometry in Protein Biomarker Research: Perspectives and Clinical Applications. *Proteomics Clin Appl* 2015, doi: 10.1002/prca.201400117, 影响因子: 2.956
43. Quantification of Lysine Acetylation and Succinylation Stoichiometry in Proteins Using Mass Spectrometric Data-Independent Acquisitions (SWATH). *J Am Soc Mass Spectr* 2016, DOI:10. 1007/s13361-016-1476-z, 影响因子: 2.869
44. Quantitative proteomics signature profiling based on network contextualization. *Biol Direct* 2015, DOI 10.1186/s13062-015-0098-x, 影响因子: 2.649

45. Bioprocess integration for human mesenchymal stem cells. *J Biotechnol* 2017, DOI: 10.1016/j.jbiotec.2017.01.014, 影响因子: 2.533
46. Comparison of protein expressions between human livers and the hepatic cell lines HepG2, Hep3B and Huh7 using SWATH and MRM-HR proteomics: Focusing on drug-metabolizing enzymes. *Drug Metab Pharmacok* 2018, doi:10.1016/j.dmpk.2018.03.003, 影响因子: 2.087
47. Proteomic Profiling of De Novo Protein Synthesis in StarvationInduced Autophagy Using Bioorthogonal Noncanonical Amino Acid Taggin. *Methods Enzymol* 2016, doi: 10.1016/bs.mie.2016.09.075, 影响因子: 1.984
48. The effect of photodamage on the female Caucasian facial stratum corneum corneome using mass spectrometry-based proteomics. *Int J Cosm Sci* 2017, DOI: 10.1111/ics.12426, 影响因子: 1.777
49. Data for mitochondrial proteomic alterations in the developing rat brain. *Data in Brief* 2014, doi: 10.1016/j.dib.2014.07.002, 影响因子: 0.341
50. Neonatal mitochondrial abnormalities due to PINK1 deficiency: Proteomics reveals early changes relevant to Parkinson's disease. *Data in Brief* 2016, doi:10.1016/j.dib.2015.11.070, 影响因子: 0.341
51. Label-free quantification of differentially expressed proteins in mouse liver cancer cells with high and low metastasis rates by a SWATH acquisition method. *Chem sci china* 2014, doi: 10.1007/s11426-014-5093-z
52. p-SRM, SWATH and HRM - Targeted Proteomics Approaches on TripleTOF 5600+ Mass Spectrometer and Their Applications in Oncology Research. *Klin Onkol* 2014, 27 (1): S110-S115
53. Label-Free Quantitation and Mapping of the ErbB2 Tumor Receptor by Multiple Protease Digestion with Data-Dependent (MS1) and Data-Independent (MS2) Acquisitions. *Int J Proteomics* 2013, doi: 10.1155/2013/791985
54. Depletion of Myofibril-Associated Proteins Using Selective Protein Extraction as a Tool in Cardiac Proteomics. *Methods Mol Biol* 2017,

doi:10.1007/7651_2017_73

55. Multi-omic network-based interrogation of rat liver metabolism following gastric bypass surgery featuring SWATH proteomics. *Technology* 2017, DOI: 10.1142/S233954781750008X
56. Chlamydospore Specific Proteins of *Candida albicans*. *Data* 2017, DOI: 10.3390/data2030026
57. THP1 cell proteome library generation using SWATH analysis. *Int J Life Sci* 2018, 6 (1):143-149
58. Genomic, Proteomic and Phenotypic Heterogeneity in HeLa Cells across Laboratories: Implications for Reproducibility of Research Results. *bioRxiv* 2018, doi: 10.1101/307

四、植物蛋白组学

1. Tomato nuclear proteome reveals the involvement of specific E2 ubiquitin-conjugating enzymes in fruit ripening. *Genome Biol* 2014, doi:10.1186/s13059-014-0548-2, 影响因子: 13.214
2. Impact of different nitrogen sources on the growth of *Arthrosira* sp. PCC 8005 under batch and continuous cultivation - A biochemical, transcriptomic and proteomic profile. *Bioresource Technology* 2017, doi: 10.1016/j.biortech.2017.03.145, 影响因子: 5.807
3. Systems-level proteomics of two ubiquitous leaf commensals reveals complementary adaptive traits for phyllosphere colonization. *Mol Cell Proteomics* 2016, M116.058164, 影响因子: 5.232
4. Analysis of dynamic protein carbonylation in rice embryo during germination through AP-SWATH. *Proteomics* 2016, doi:10.1002/pmic.201500248, 影响因子: 3.532
5. Extensive mass spectrometry proteomics data of *Persicaria minor* herb

- upon methyl jasmonate treatment. Data in Brief 2018, doi: 10.1016/j.dib.2017.09.063, 影响因子：0.341
6. Proteomic analysis of pitcher fluid from Nepenthes. Data in Brief 2018, doi: org/10.1016/j.dib.2018.01.037, 影响因子：0.341

五、动物蛋白组学

1. Proteome-wide association studies identify biochemical modules associated with a wing-size phenotype in *Drosophila melanogaster*. Nat Commun 2016, DOI: 10.1038/ncomms12649, 影响因子：12.124
2. Integrated Transcriptomic and Proteomic Analyses Suggest the Participation of Endogenous Protease Inhibitors in the Regulation of Protease Gene Expression in *Helicoverpa armigera*. Mol Cell Proteomics 2018, doi: 10.1074/mcp.RA117.000533, 影响因子：5.232
3. Discrimination of dried sea cucumber (*Apostichopus japonicus*) products from different geographical origins by sequential windowed acquisition of all theoretical fragment ion mass spectra (SWATH-MS)-based proteomic analysis and chemometrics. Food Chem 2018, doi:10.1016/j.food-chem.2018.08.082, 影响因子：4.946
4. Label-free Quantitative Analysis of Changes in Broiler Liver Proteins under Heat Stress using SWATH-MS Technology. Sci Rep 2015, DOI: 10.1038/srep15119, 影响因子：4.122
5. The Pig PeptideAtlas: A resource for systems biology in animal production and biomedicine. Proteomics 2016, doi:0.1002/pmic.201500195, 影响因子：3.532
6. SWATH-based quantitative proteomics reveals the mechanism of enhanced *Bombyx mori* nucleopolyhedrovirus-resistance in silkworm reared on UV-B treated mulberry leaves. Proteomics 2017, doi:10.1002/pmic.201600383, 影响因子：3.532

7. Extensive dataset of boar seminal plasma proteome displaying putative reproductive functions of identified proteins. Data in Brief 2016, doi:10.1016/j.jprot.2016.04.026, 影响因子：0.341
8. SWATH-MS data of Drosophila melanogaster proteome dynamics during embryogenesis. Data in Brief 2016, doi:10.1016/j.dib.2016.10.009, 影响因子：0.341
9. Definition of a RACK1 interaction network in Drosophila melanogaster using SWATH-MS. G3 (Bethesda) 2017, doi: 10.1534/g3.117.042564

六、微生物蛋白组学（发酵）

1. Absolute Proteome Composition and Dynamics during Dormancy and Resuscitation of Mycobacterium tuberculosis. Cell Host Microbe 2015, doi:10.1016/j.chom.2015.06.001, 影响因子：14.946
2. Mass Spectral Enhanced Detection of Ubis Using SWATH Acquisition: MEDUSA-Simultaneous Quantification of SUMO and Ubiquitin-Derived Isopeptides. J Am Chem Soc 2014, DOI: 10.1007/s13361-014-0835-x, 影响因子：13.858
3. Reproducible and Consistent Quantification of the *Saccharomyces cerevisiae* Proteome by SWATH-mass spectrometry. Mol Cell Proteomics 2015, doi:10.1074/mcp.M113.035550, 影响因子：5.232
4. Daily rhythms in the cyanobacterium *synechococcus elongatus* probed by high-resolution mass spectrometry-based proteomics reveals a small defined set of cyclic proteins. Mol Cell Proteomics 2014, doi: 10.1074/mcp.M113.035840, 影响因子：5.232
5. Simultaneous quantification of viral antigen expression kinetics using data-independent mass spectrometry. Mol Cell Proteomics 2015, doi: 10.1074/mcp.M114.047373, 影响因子：5.232

6. Effects of lipid-lowering drugs on vancomycin susceptibility of mycobacteria. *Antimicrob Agents Chemother* 2016, doi:10.1128/AAC.00872-16, 影响因子：4.255
7. Response induced in *Mycoplasma gallisepticum* under heat shock might be relevant to infection process. *Sci Rep* 2017, DOI:10.1038/s41598-017-09237-7, 影响因子：4.122
8. Understanding the dimorphic lifestyles of human gastric pathogen *Helicobacter pylori* using the SWATH-based proteomics approach. *Sci Rep* 2016, DOI: 10.1038/srep26784, 影响因子：4.122
9. Selective enrichment of mycobacterial proteins from infected host macrophages. *Sci Rep* 2015, DOI: 10.1038/srep13430, 影响因子：4.122
10. SWATH label-free proteomics analyses revealed the roles of oxidative stress and antioxidant defensing system in sclerotia formation of *Polyporus umbellatus*. *Sci Rep* 2017, DOI: 10.1038/srep41283, 影响因子：4.122
11. The use of SWATH to analyse the dynamic changes of bacterial proteome of carbapanemase-producing *Escherichia coli* under antibiotic pressure. *Sci Rep* 2018, DOI:10.1038/s41598-018-21984-9, 影响因子：4.122
12. Glycation inhibitors extend yeast chronological lifespan by reducing advanced glycation end products and by back regulation of proteins involved in mitochondrial respiration. *J Proteomics* 2017, doi:10.1016/j.jprot.2017.01.015, 影响因子：3.722
13. *Thermoanaerobacter thermohydrosulfuricus WC1 Shows Protein Complement Stability during Fermentation of Key Lignocellulose-Derived Substrates*. *Appl Environ Microb* 2014, doi:10.1128/AEM.03555-13, 影响因子：3.633
14. In-depth comparative proteomic analysis of yeast proteome using iTRAQ and SWATH based MS. *Mol BioSyst* 2015, DOI: 10.1039/c5mb00234f , 影响因子：2.759

15. Proteomic Analysis of Alterations in *Aeromonas hydrophila* Outer Membrane Proteins in Response to Oxytetracycline Stress. *Microb Drug Resist* 2018, doi:10.1089/mdr.2017.0324, 影响因子: 2.344
16. Proteomic Analysis of *Vibrio parahaemolyticus* Under Cold Stress. *Curr Microbiol* 2017, doi:10.1007/s00284-017-1345-4, 影响因子: 1.373
17. SWATH-based Comparative Proteomic Analysis of the *Mycobacterium bovis*. *Bull Korean Chem Soc* 2014, doi:10.5012/bkcs.2014.35.3.933, 影响因子: 0.522
18. Chlamydospore Specific Proteins of *Candida albicans*. *Data* 2017, DOI: 10.3390/data2030026

七、蛋白相互作用、食品组学、生命科学研究等蛋白组学应用

1. Mapping differential interactomes by affinity purification coupled with data independent mass spectrometry acquisition. *Nat Med* 2013, doi:10.1038/nmeth.2702, 影响因子: 32.621
2. Interaction Domains of Sos1/Grb2 Are Finely Tuned for Cooperative Control of Embryonic Stem Cell Fate. *Cell* 2013, doi: 10.1016/j.cell.2013.01.056, 影响因子: 30.41
3. Quantifying protein interaction dynamics by SWATH mass spectrometry: Application to the 14-3-3 system. *Nature method* 2013, DOI: 10.1038/nmeth.2703, 影响因子: 29.886
4. Targeted proteomics *Nature method* 2013 VOL.10 NO.1 | JANUARY 2013 29.886 Methionine Metabolism Alters Oxidative Stress Resistance via the Pentose Phosphate Pathway. *Antioxid Redox Sig* 2016, DOI: 10.1089/ars.2015.6516, 影响因子: 6.53
5. Pyruvate kinase is a dosage-dependent regulator of cellular amino acid homeostasis. *Oncotarget* 2012, doi: 10.18632/oncotarget.730, 影响因

子：5.168

6. A modular and adaptive mass spectrometry-based platform for support of bioprocess development toward optimal host cell protein clearance. *mABs* 2017, doi:10.1080/19420862.2017.1303023, 影响因子：5.165
7. A multiplex and label-free relative quantification approach for studying protein abundance of drug metabolizing enzymes in human liver microsomes using SWATH-MS. *J Proteome Res* 2017, DOI: 10.1021/acs.jproteome.7b00505, 影响因子：3.95
8. SWATH mass spectrometry as a tool for quantitative profiling of the matriosome. *J Proteomics* 2018, doi:10.1016/j.jprot.2018.02.026, 影响因子：3.722
9. Protein tyrosine adduct in humans self-poisoned by chlorpyrifos. *Toxicol Appl Pharmacol* 2013, doi: 10.1016/j.taap.2013.03.021, 影响因子：3.616
10. Analysis of protein isoforms: can we do it better. *Proteomics* 2012, doi:10.1002/pmic.201200161, 影响因子：3.532
11. Mass spectrometric protein maps for biomarker discovery and clinical research. *Expert Rev Mol Diagn* 2013, doi:10.1586/14737159.2013.845089, 影响因子：3.326
12. Human Tissue Distribution of Carbonyl Reductase 1 Using Proteomic Approach With Liquid Chromatography-Tandem Mass Spectrometry. *J Pharm Sci* 2017, doi:10.1016/j.xphs.2017.01.024, 影响因子：3.075
13. Regulation of highly homologous major urinary proteins in house mice quantified with label-free proteomic methods. *Mol BioSyst* 2016, DOI: 10.1039/c6mb00278a, 影响因子：2.759
14. HX-MS2 for high performance conformational analysis of complex protein states. *Protein Sci* 2015, DOI: 10.1002/pro.2707, 影响因子：2.41
15. Drug Target Identification Using an iTRAQ-Based Quantitative Chemical

Proteomics Approach— Based on a Target Profiling Study of Andrographolide Methods. Enzymol 2016, doi:10.1016/bs.mie.2016.09.049, 影响因子：1.984

16. Authentication of leguminous-based products by targeted biomarkers using high resolution time of flight mass spectrometry. LWT - Food Sci Technol 2017, doi: 10.1016/j.lwt.2017.12.034, 影响因子：3.129
17. Resource misallocation as a mediator of fitness costs in antibiotic resistance. bioRxiv 2018, doi: 10.1101/456434

代谢组学

1. MS-DIAL: data-independent MS/MS deconvolution for comprehensive metabolome analysis. *Nature method* 2015, doi:10.1038/nmeth.3393, 影响因子：29.886
2. Metabolic Reaction Network-based Recursive Metabolite Identification for Untargeted Metabolomics. *Nat Commun* 2019, doi: 10.1038/s41467-019-09550-x, 影响因子：12.144
3. Integrated Quantification and Identification of Aldehydes and Ketones in Biological Samples. *Anal Chem* 2014, doi:10.1021/ac500810r, 影响因子：6.32
4. Discovering Regulated Metabolite Families in Untargeted Metabolomics Studies. *Anal Chem* 2016, DOI: 10.1021/acs.analchem.6b01569, 影响因子：6.32
5. MetDIA: Targeted Metabolite Extraction of Multiplexed MS/MS Spectra Generated by Data-Independent Acquisition. *Anal Chem* 2016, DOI: 10.1021/acs.analchem.6b02122, 影响因子：6.32
6. Customized consensus spectral library building for untargeted quantitative metabolomics analysis using data independent acquisition mass spectrometry and MetaboDIA workflow. *Anal Chem* 2017, DOI: 10.1021/acs.analchem.6b05006, 影响因子：6.32
7. Ion-pair selection method for pseudotargeted metabolomics based on SWATH MS acquisition and its application in differential metabolite discovery of Type 2 diabetes. *Anal Chem* 2018, DOI: 10.1021/acs.analchem.8b02377, 影响因子：6.32
8. Ultrafast polyphenol metabolomics of red wines using microLC-MS/MS. *J Agric Food Chem* 2016, DOI: 10.1021/acs.jafc.5b04890 , 影响因子：3.412
9. SWATH acquisition mode for drug metabolism and metabolomics investiga-

- tions. *Bioanalysis* 2016, DOI: 10.4155/bio-2016-0141, 影响因子: 2.67
10. SWATH-Based Metabolomics of Follicular Fluid in Patients Shows That Progesterone Adversely Affects Oocyte Quality. *Biomed Res Int* 2018, doi: 10.1155/2018/1780391, 影响因子: 2.583
11. SWATH data independent acquisition mass spectrometry for metabolomics. *TrAC-Trend Anal Chem* 2018, doi:10.1016/j.trac.2018.10.014, 影响因子: 7.034

脂质组学

1. Novel approach in LC-MS/MS using MRM to generate a full profile of acyl-CoAs: discovery of acyl-dephospho-CoAs. *J Lipid Res* 2014, doi: 10.1194/jlr.D045112, 影响因子: 4.81
2. Absolute quantitative lipidomics reveals lipidome-wide alterations in aging brain. *Metabolomics* 2018, doi:10.1007/s11306-017-1304-x, 影响因子: 3.511
3. Early lipid changes in acute kidney injury using SWATH lipidomics coupled with MALDI tissue imaging. *Am J Physiol Ren Physiol* 2016, DOI: 10.1152/ajprenal.00100.2016, 影响因子: 3.164
4. Comprehensive MS/MS Profiling by UHPLC-ESI-QTOF-MS/MS using SWATH Data-independent Acquisition for the Study of Platelet Lipidomes in Coronary Artery Disease. *Anal Chim Acta* 2018, doi:10.1016/j.aca.2018.08.060 , 影响因子: 5.123
5. Comparison of simple monophasic versus classical biphasic extraction protocols for comprehensive UHPLC-MS/MS lipidomic analysis of Hela cells. *Anal Chim Acta* 2018, DOI:10.1016/j.aca.2018.10.035, 影响因子: 5.123
6. Comparative Lipidomics of *Caenorhabditis elegans* Metabolic Disease Models by SWATH Non-Targeted Tandem Mass Spectrometry. *Metabolites* 2015, doi: 10.3390/metabo5040677

药物分析

1. Comparison of Information-Dependent Acquisition, SWATH, and MSAll Techniques in Metabolite Identification Study Employing Ultrahigh-Performance Liquid Chromatography-Quadrupole Time of-Flight Mass Spectrometry. *Anal Chem* 2014, doi: 10.1021/ac403385y, 影响因子: 6.32
2. High-resolution mass spectrometry for integrated qualitative and quantitative analysis of pharmaceuticals in biological matrices. *Anal Bioanal Chem* 2012, DOI 10.1007/s00216-011-5641-8, 影响因子: 3.2
3. Nontargeted SWATH acquisition mode for metabolites identification of osthole in rats using ultra-high-performance liquid chromatography coupled to quadrupole time-of-flight mass spectrometry. *RSC Adv* 2018, DOI:10.1039/c8ra01221k , 影响因子: 2.936
4. A practical strategy for the characterization of ponicidin metabolites in vivo and in vitro by UHPLC-Q-TOF-MS based on nontargeted SWATH data acquisition. *J Pharma Biomed Anal* 2017, doi:10.1016/j.jpba.2017.08.003, 影响因子: 2.831
5. Simultaneous quantitative analysis of polyethylene glycol (PEG), PEGylated paclitaxel and paclitaxel in rats by MS/MSALL technique with hybrid quadrupole time-of-flight mass spectrometry. *J Pharma Biomed Anal* 2017, doi:10.1016/j.jpba.2017.06.053, 影响因子: 2.831
6. Quantification of acetaminophen and two of its metabolites in human plasma by ultra-high performance liquid chromatography-low and high resolution tandem mass spectrometry. *J Chromatogr B* 2012, doi:10.1016/j.jchromb.2012.07.009, 影响因子: 2.441

毒物、污染物分析

1. Liquid Chromatography, in Combination with a Quadrupole Time-of-Flight Instrument (LC QTOF), with Sequential Window Acquisition of All Theoretical Fragment-Ion Spectra (SWATH) Acquisition: Systematic Studies on Its Use for Screenings in Clinical and Forensic Toxicology and Comparison with Information-Dependent Acquisition (IDA). *Anal Chem* 2014, doi:10.1021/ac503144p, 影响因子：6.32
2. In vitro and in vivo human metabolism of a new synthetic cannabinoid NM-2201 (CBL-2201). *Forensic Toxicol* 2017, DOI: 10.1007/s11419-016-0326-9, 影响因子：3.924
3. Nontargeted SWATH acquisition for identifying 47 synthetic cannabinoid metabolites in human urine by liquid chromatography-high-resolution tandem mass spectrometry. *Anal Bioanal Chem* 2014, DOI 10.1007/s00216-014-8118-8, 影响因子：3.2
4. Applying ‘Sequential Windowed Acquisition of All Theoretical Fragment Ion Mass Spectra’ (SWATH) for systematic toxicological analysis with liquid chromatography-high-resolution tandem mass spectrometry. *Anal Bioanal Chem* 2014, DOI <https://doi.org/10.1007/s00216-014-8262-1>, 影响因子：3.2
5. In vitro metabolism of the synthetic cannabinoids CUMYL-PINACA, 5F-CUMYL-PINACA, CUMYL-4CN-BINACA, 5F-CUMYL-P7AICA and CUMYL-4CN-B7AICA. *Drug Test Anal* 2017, DOI: 10.1002/dta.2298, 影响因子：2.993
6. Metabolism of novel opioid agonists U-47700 and U-49900 using human liver microsomes with confirmation in authentic urine specimens from drug users. *Drug Test Anal* 2018, DOI: 10.1002/dta.2228, 影响因子：2.99
7. Parameter Optimization for Feature and Hit Generation in a General Unknown Screening Method - Proof of Concept Study Using a Design of

Experiment Approach for a High Resolution Mass Spectrometry Procedure after Data Independent Acquisition. Anal Chem 2018, DOI: 10.1021/acs.analchem.7b05387, 影响因子: 6.32

8. Ozonation, photocatalysis and photocatalytic ozonation of diuron. Intermediates identification. Chem Eng J 2016, doi:10.1016/j.cej.2016.02.005 , 影响因子: 6.216
9. High-resolution mass spectrometry with data independent acquisition for the comprehensive nontargeted analysis of migrating chemicals coming from multilayer plastic packaging materials used for fruit purée and juice. Talanta 2018, doi:10.1016/j.talanta.2018.08.023 , 影响因子: 4.244

您的成功就是我们工作的动力 我们视为己任。

作为SCIEX的客户，您将得到世界一流的售后支持，无论在何地，我们都将是您最为可靠的伙伴，为您解决难题，提供方案以及最大化提高工作效率。

我们的售后工程师提供全线LC-MS/MS系统支持。无论您的离子源，自动进样器或者实际样品分析需要帮助，您都将得到及时反馈。我们将最大努力保证仪器的正常使用，帮助您完成科研工作。

我们的应用科学家提供可靠的工作流程，帮助您简化样品制备，减少人工操作步骤。帮助您开发方法以提高分析通量。通过提供简单易用的iMethod™方法，帮助您快速开展工作。我们也可以通过电话实现远程支持。

我们的资深专家为您量身订做适合您实验室的培训课程，提高工作效率。您可以到我们的Demo Lab参加LC-MS/MS技术培训及应用课程，也可以通过网络 e-learning工具进行学习。

我们的售后支持团队随时为您提供最新产品信息，软件更新，分析方法及仪器维护，确保您在竞争中独占鳌头。

我们将竭尽所能为您提供高效优质的服务。

更多信息，请访问：www.sciex.com.cn

www.sciex.com



For Research Use Only. Not for use in Diagnostics Procedures.

AB Sciex is operating as SCIEX.

© 2019. AB Sciex. The trademarks mentioned herein are the property of AB Sciex Pte.

Ltd. or their respective owners. AB SCIEX™ is being used under license.

RUO-MKT-07-9450-ZH-A



SCIEX中国公司

北京分公司

地址：北京市朝阳区酒仙桥中路24号院
1号楼5层

电话：010-58081388
传真：010-58081390

全国免费垂询电话：800 820 3488, 400 821 3897

SCIEXNow™服务热线：800 820 3488, 400 821 3897

上海公司及亚太区应用支持中心

地址：上海市长宁区福泉北路518号
1座502室

电话：021-24197200
传真：021-24197333

网址：sciex.com.cn

广州分公司

地址：广州市天河区珠江西路15号
珠江城1907室

电话：020-85100200
传真：020-38760835

微博：@SCIEX

服务邮箱：Service.china@sciex.com